

D-MAVT, D-MATL

Mock Exam Analysis III

401-0363-10L

Last Name

XX

First Name

XX

Legi-Nr.

XX-000-000

Exam-No.

000

Please do not turn the page yet!

Please take note of the information on the answer-booklet.

Laplace Transforms: $(F = \mathcal{L}(f))$

	$f(t)$	$F(s)$		$f(t)$	$F(s)$		$f(t)$	$F(s)$
1)	1	$\frac{1}{s}$	5)	$t^a, a > 0$	$\frac{\Gamma(a+1)}{s^{a+1}}$	9)	$\cosh(at)$	$\frac{s}{s^2 - a^2}$
2)	t	$\frac{1}{s^2}$	6)	e^{at}	$\frac{1}{s-a}$	10)	$\sinh(at)$	$\frac{a}{s^2 - a^2}$
3)	t^2	$\frac{2}{s^3}$	7)	$\cos(\omega t)$	$\frac{s}{s^2 + \omega^2}$	11)	$u(t-a)g(t-a)$	$\mathcal{L}(g)e^{-as}$
4)	$t^n, n \in \mathbb{Z}_{\geq 0}$	$\frac{n!}{s^{n+1}}$	8)	$\sin(\omega t)$	$\frac{\omega}{s^2 + \omega^2}$	12)	$\delta(t-a)$	e^{-as}

(Γ = Gamma function, u = Heaviside function, δ = Dirac's delta function)

Fourier transforms:

	$f(x)$	$\widehat{f}(\omega)$		$f(x)$	$\widehat{f}(\omega)$		$f(x)$	$\widehat{f}(\omega)$
1)	e^{-ax^2}	$\frac{1}{\sqrt{2a}}e^{\frac{-\omega^2}{4a}}$	2)	$\begin{cases} e^{-ax}, & x \geq 0, \\ 0, & x < 0. \end{cases}$	$\frac{1}{\sqrt{2\pi}(a+i\omega)}$	3)	$\begin{cases} 1, & x < 1, \\ 0, & x > 1. \end{cases}$	$\sqrt{\frac{2}{\pi}} \frac{\sin(\omega)}{\omega}$

Indefinite Integrals: $(n \in \mathbb{Z}_{\geq 1})$

1)	$\int x \cos\left(\frac{n\pi}{L}x\right) dx = \frac{\cos\left(\frac{n\pi}{L}x\right) + \left(\frac{n\pi}{L}\right)x \sin\left(\frac{n\pi}{L}x\right)}{\left(\frac{n\pi}{L}\right)^2} \quad (+\text{constant})$
2)	$\int x^2 \cos\left(\frac{n\pi}{L}x\right) dx = \frac{\left(\left(\frac{n\pi}{L}\right)^2 x^2 - 2\right) \sin\left(\frac{n\pi}{L}x\right) + 2\left(\frac{n\pi}{L}\right)x \cos\left(\frac{n\pi}{L}x\right)}{\left(\frac{n\pi}{L}\right)^3} \quad (+\text{constant})$
3)	$\int x \sin\left(\frac{n\pi}{L}x\right) dx = \frac{\sin\left(\frac{n\pi}{L}x\right) - \left(\frac{n\pi}{L}\right)x \cos\left(\frac{n\pi}{L}x\right)}{\left(\frac{n\pi}{L}\right)^2} \quad (+\text{constant})$
4)	$\int x^2 \sin\left(\frac{n\pi}{L}x\right) dx = \frac{\left(2 - \left(\frac{n\pi}{L}\right)^2 x^2\right) \cos\left(\frac{n\pi}{L}x\right) + 2\left(\frac{n\pi}{L}\right)x \sin\left(\frac{n\pi}{L}x\right)}{\left(\frac{n\pi}{L}\right)^3} \quad (+\text{constant})$
5)	$\int \frac{1}{1+x^2} dx = \arctan(x) \quad (+\text{constant})$

Question 1

1.MC1 [3 Points] Let f be a solution of the following ordinary differential equation (ODE),

$$\begin{cases} f''(t) + \omega^2 f(t) = 0, & t > 0 \\ f(0) = 1, & f'(0) = 2\omega, \end{cases}$$

where $\omega > 0$ is a positive constant. Find the Laplace transform $\mathcal{L}(f) = F$ of the function f .

- (A) $F(s) = \frac{s}{s^2 + \omega^2} + \frac{2\omega}{s^2 + \omega^2}$.
- (B) $F(s) = \frac{1}{s^2 + \omega^2} + \frac{2s\omega}{s^2 + \omega^2}$.
- (C) $F(s) = \frac{2\omega}{s + \omega^2}$.
- (D) $F(s) = \frac{2\omega}{s^2 + \omega}$.

1.MC2 [3 Points] Find the inverse Laplace transform of the following function

$$F(s) = \frac{s + 2}{s^2 - 10s + 25}.$$

- (A) $f(t) = e^{-5t}(1 + 7t)$.
- (B) $f(t) = e^{5t}(1 + 7t)$.
- (C) $f(t) = e^{5t}(1 - 7t)$.
- (D) $f(t) = e^{-5t}(1 - 7t)$.

1.MC3 [3 Points] Let f be a continuous function such that $\lim_{x \rightarrow \infty} f(x) = 0$. Solve the following differential equation using the Fourier transform

$$f(x) + f'(x) + 4f''(x) = \sqrt{2\pi}e^{-\pi x^2}.$$

- (A) $f(x) = \int_{-\infty}^{\infty} \frac{1}{1+i\omega-4\omega^2} e^{-\frac{\omega^2}{4\pi}} e^{-i\omega x} d\omega$.
- (B) $f(x) = \int_{-\infty}^{\infty} \frac{1}{1+i\omega+4\omega^2} e^{-\frac{\omega^2}{4\pi}} e^{i\omega x} d\omega$.
- (C) $f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{1+i\omega-4\omega^2} e^{-\frac{\omega^2}{4\pi}} e^{i\omega x} d\omega$.
- (D) $f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{1+i\omega+4\omega^2} e^{-\frac{\omega^2}{4\pi}} e^{-i\omega x} d\omega$.

1.MC4 [3 Points] The complex Fourier series of the function $\cosh(ax)$ on the interval $[-\pi, \pi)$ is given by

$$\sum_{n=-\infty}^{+\infty} \frac{(-1)^n a \sinh(a\pi)}{\pi(n^2 + a^2)} e^{inx}.$$

Find the value of the numerical series

$$\sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2}$$

- (A) $\sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2} = \frac{\pi}{2a \sinh(a\pi)}.$
(B) $\sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2} = \frac{\pi}{2a \sinh(a\pi)} - \frac{1}{2a^2}.$
(C) $\sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2} = \frac{2\pi}{a \sinh(a\pi)} - \frac{2}{a^2}.$
(D) $\sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2} = \frac{2\pi}{a \sinh(a\pi)}.$

1.MC5 [3 Points] Determine if the following function is even, odd, or neither and if it is periodic or not. If the function is periodic, determine the fundamental period.

$$15 \cos(3x) + 3 \cos(4x).$$

- (A) The function is even and periodic of fundamental period 2π .
(B) The function is odd and periodic of fundamental period 2π .
(C) The function is even and periodic of fundamental period π .
(D) The function is odd and periodic of fundamental period π .

1.MC6 [3 Points] Consider the following PDE (partial differential equation) for the function $u = u(x, y)$:

$$4u_{xx} + xu_x + 6u_{xy} - yu_y = -7u_{yy} + u_x.$$

Is the PDE hyperbolic, parabolic, elliptic or of mixed type ?

- (A) hyperbolic.
(B) elliptic.
(C) parabolic.
(D) mixed type.

1.MC7 [3 Points] Wave equation with D'Alembert solution.

Let $u(x, t)$ be the solution of the following problem

$$\begin{cases} u_{tt} = u_{xx}, & x \in \mathbb{R}, t > 0, \\ u(x, 0) = f(x) = \begin{cases} 2, & |x| \leq 14 \\ 0, & |x| > 14 \end{cases} & x \in \mathbb{R}, \\ u_t(x, 0) = g(x) = \begin{cases} 1, & 3 \leq x \leq 4 \\ 0, & x \notin [3, 4] \end{cases} & x \in \mathbb{R}. \end{cases}$$

Find the values of u at the point $(x, t) = (10, 7)$, i.e. find $u(10, 7)$

- (A) $u(10, 7) = 8$.
- (B) $u(10, 7) = 10$.
- (C) $u(10, 7) = \frac{3}{2}$.
- (D) $u(10, 7) = \frac{5}{2}$.

1.MC8 [3 Points] Let $u = u(x, y)$ be a harmonic function in D_2 the disk of radius 2 centred at 0.

The maximum value of u is at $(x, y) = (\sqrt{2}, -\sqrt{2})$, i.e. $\max_{D_2} u(x, y) = u(\sqrt{2}, -\sqrt{2})$.

Which of the following statements is true?

- (A) u is not constant in D_2 .
- (B) u is constant in D_2 .
- (C) There exists another point (x', y') in D_2 such that $u(x, y) = u(x', y')$.
- (D) We cannot conclude that (A), (B) and (C) are true for every u .

1.MC9 [3 Points] Consider the Neumann problem for the following PDE,

$$\begin{cases} \nabla^2 u = f, & \text{in } D_2, \\ \frac{\partial u}{\partial n} = g, & \text{on } \partial D_2, \end{cases}$$

with D_2 the disk of radius 2 centred at 0 and f and g are two given functions such that

$$\int_{D_2} f(x) \, dx = 2, \quad \text{and} \quad \int_{\partial D_2} g(x) \, dx = 2.$$

Which of the following is true:

- (A) There are infinitely many solutions.
- (B) There is no solution.
- (C) There are two solutions.
- (D) We cannot conclude that (A), (B), or (C) are true.

Question 2

2.Q1 [15 Points] Separation of variables for the Heat equation

Consider the following time-dependent version of the Heat equation on the interval $[0, 1]$. We also impose boundary conditions and we look for a solution $u = u(x, t)$ such that:

$$\begin{cases} u_t(x, t) = a(t)u_{xx}(x, t), & x \in [0, 1], t \in [0, +\infty), \\ u(0, t) = 0, & t \in [0, +\infty), \\ u(1, t) = 0, & t \in [0, +\infty), \\ u(x, 0) = 2 \sin(3\pi x) + \sin(7\pi x), & x \in [0, 1], \end{cases}$$

where $a(t)$ is a given continuous function. Find the solution $u(x, t)$ using separation of variable. Proceed as in the lecture and adapt the steps if necessary.

Question 3

3.Q1 [10 Points] Dirichlet problem on a region with symmetries

Find the solution $u(r, \theta)$ of the following Dirichlet problem on the disk of radius R in polar coordinates:

$$\begin{cases} \nabla^2 u = 0, & 0 \leq r \leq R, 0 \leq \theta \leq 2\pi, \\ u(R, \theta) = \sin^2(\theta) + 8 \cos^3(\theta), & 0 \leq \theta \leq 2\pi. \end{cases}$$

You should give the answer without unsolved integral and you can use the formulas developed in the lecture.

[Hint: Don't try to find the solution in the Poisson integral form.]

[Hint: Remember the trigonometric formulas

$$\sin^2(\theta) = \frac{1}{2} - \frac{1}{2} \cos(2\theta) \quad \text{and} \quad \cos^3(\theta) = \frac{3}{4} \cos(\theta) + \frac{1}{4} \cos(3\theta).]$$