

Problems and suggested solution

Laplace Transforms: ($F = \mathcal{L}(f)$) (u = Heaviside function, δ = Dirac's delta function)

	$f(t)$	$F(s)$		$f(t)$	$F(s)$		$f(t)$	$F(s)$
1)	1	$\frac{1}{s}$	5)	$t^a, a > 0$	$\frac{\Gamma(a+1)}{s^{a+1}}$	9)	$\cosh(at)$	$\frac{s}{s^2-a^2}$
2)	t	$\frac{1}{s^2}$	6)	e^{at}	$\frac{1}{s-a}$	10)	$\sinh(at)$	$\frac{a}{s^2-a^2}$
3)	t^2	$\frac{2}{s^3}$	7)	$\cos(\omega t)$	$\frac{s}{s^2+\omega^2}$	11)	$u(t-a)g(t-a)$	$\mathcal{L}(g)e^{-as}$
4)	$t^n, n \in \mathbb{Z}_{\geq 0}$	$\frac{n!}{s^{n+1}}$	8)	$\sin(\omega t)$	$\frac{\omega}{s^2+\omega^2}$	12)	$\delta(t-a)$	e^{-as}

Fourier transforms:

	$f(x)$	$\hat{f}(\omega)$		$f(x)$	$\hat{f}(\omega)$		$f(x)$	$\hat{f}(\omega)$
1)	e^{-ax^2}	$\frac{1}{\sqrt{2a}}e^{-\frac{\omega^2}{4a}}$	2)	$\begin{cases} e^{-ax}, & x \geq 0, \\ 0, & x < 0. \end{cases}$	$\frac{1}{\sqrt{2\pi}(a+i\omega)}$	3)	$\begin{cases} 1, & x < 1, \\ 0, & x > 1. \end{cases}$	$\sqrt{\frac{2}{\pi}} \frac{\sin(\omega)}{\omega}$

Indefinite Integrals: ($n \in \mathbb{Z}_{\geq 1}$)

1)	$\int x \cos\left(\frac{n\pi}{L}x\right) dx = \frac{\cos\left(\frac{n\pi}{L}x\right) + \left(\frac{n\pi}{L}\right)x \sin\left(\frac{n\pi}{L}x\right)}{\left(\frac{n\pi}{L}\right)^2} \quad (+\text{constant})$
2)	$\int x^2 \cos\left(\frac{n\pi}{L}x\right) dx = \frac{\left(\left(\frac{n\pi}{L}\right)^2 x^2 - 2\right) \sin\left(\frac{n\pi}{L}x\right) + 2\left(\frac{n\pi}{L}\right)x \cos\left(\frac{n\pi}{L}x\right)}{\left(\frac{n\pi}{L}\right)^3} \quad (+\text{constant})$
3)	$\int x \sin\left(\frac{n\pi}{L}x\right) dx = \frac{\sin\left(\frac{n\pi}{L}x\right) - \left(\frac{n\pi}{L}\right)x \cos\left(\frac{n\pi}{L}x\right)}{\left(\frac{n\pi}{L}\right)^2} \quad (+\text{constant})$
4)	$\int x^2 \sin\left(\frac{n\pi}{L}x\right) dx = \frac{\left(2 - \left(\frac{n\pi}{L}\right)^2 x^2\right) \cos\left(\frac{n\pi}{L}x\right) + 2\left(\frac{n\pi}{L}\right)x \sin\left(\frac{n\pi}{L}x\right)}{\left(\frac{n\pi}{L}\right)^3} \quad (+\text{constant})$
5)	$\int \frac{1}{1+x^2} dx = \arctan(x) \quad (+\text{constant})$

You can use these formulas without justification.

Question 1

1.MC1 [3 Points] Let f be a solution of the following ordinary differential equation (ODE),

$$\begin{cases} f''(t) + \omega^2 f(t) = 0, & t > 0 \\ f(0) = 1, & f'(0) = 2\omega, \end{cases}$$

where $\omega > 0$ is a positive constant. Find the Laplace transform $\mathcal{L}(f) = F$ of the function f .

- (A) $F(s) = \frac{s}{s^2 + \omega^2} + \frac{2\omega}{s^2 + \omega^2}$.
 (B) $F(s) = \frac{1}{s^2 + \omega^2} + \frac{2s\omega}{s^2 + \omega^2}$.
 (C) $F(s) = \frac{2\omega}{s + \omega^2}$.
 (D) $F(s) = \frac{2\omega}{s^2 + \omega}$.

Solution:

The solution is (A) $F(s) = \frac{s}{s^2 + \omega^2} + \frac{2\omega}{s^2 + \omega^2}$.

We apply the Laplace transform to the ODE in the initial value problem. We denote by $F = \mathcal{L}(f)$ the Laplace transform of the function f , and we denote the variable in the new domain by s as usual (so $F = F(s)$).

The first term to transform is the second derivative f'' , for which we use the formula:

$$\mathcal{L}(f'') = s^2 F - s f(0) - f'(0) = s^2 F - s - 2\omega.$$

Then we have $\mathcal{L}(\omega^2 f) = \omega^2 F$ (by linearity). In conclusion the ODE becomes the following algebraic equation:

$$s^2 F - s - 2\omega + \omega^2 F = 0 \quad \implies \quad F = \frac{s}{s^2 + \omega^2} + \frac{2\omega}{s^2 + \omega^2}.$$

1.MC2 [3 Points] Find the inverse Laplace transform of the following function

$$F(s) = \frac{s + 2}{s^2 - 10s + 25}.$$

- (A) $f(t) = e^{-5t}(1 + 7t)$.
 (B) $f(t) = e^{5t}(1 + 7t)$.
 (C) $f(t) = e^{5t}(1 - 7t)$.
 (D) $f(t) = e^{-5t}(1 - 7t)$.

Solution:

The solution is (B) $f(t) = e^{5t}(1 + 7t)$.

We have

$$F(s) = \frac{s + 2}{s^2 - 10s + 25} = \frac{s + 2}{(s - 5)^2} = \frac{(s - 5) + 7}{(s - 5)^2} = \frac{1}{s - 5} + \frac{7}{(s - 5)^2}.$$

Therefore,

$$f(t) = \mathcal{L}^{-1} \left(\frac{1}{s-5} + \frac{7}{(s-5)^2} \right) = \mathcal{L}^{-1} \left(\frac{1}{s-5} \right) + \mathcal{L}^{-1} \left(\frac{7}{(s-5)^2} \right) = e^{5t} + e^{5t} 7t = e^{5t}(1 + 7t).$$

1.MC3 [3 Points] Let f be a continuous function such that $\lim_{x \rightarrow \infty} f(x) = 0$. Solve the following differential equation using the Fourier transform

$$f(x) + f'(x) + 4f''(x) = \sqrt{2\pi} e^{-\pi x^2}.$$

- (A) $f(x) = \int_{-\infty}^{\infty} \frac{1}{1+i\omega-4\omega^2} e^{-\frac{\omega^2}{4\pi}} e^{-i\omega x} d\omega.$
- (B) $f(x) = \int_{-\infty}^{\infty} \frac{1}{1+i\omega+4\omega^2} e^{-\frac{\omega^2}{4\pi}} e^{i\omega x} d\omega.$
- (C) $f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{1+i\omega-4\omega^2} e^{-\frac{\omega^2}{4\pi}} e^{i\omega x} d\omega.$
- (D) $f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{1+i\omega+4\omega^2} e^{-\frac{\omega^2}{4\pi}} e^{-i\omega x} d\omega.$

Solution:

The solution is (C) $f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{1+i\omega-4\omega^2} e^{-\frac{\omega^2}{4\pi}} e^{i\omega x} d\omega.$

We take the Fourier transform on both sides of the equation. (We use the table above and the property of the Fourier transform for a derivative.)

$$\hat{f}(\omega) + i\omega \hat{f}(\omega) - 4\omega^2 \hat{f}(\omega) = \sqrt{2\pi} \frac{1}{\sqrt{2\pi}} e^{-\frac{\omega^2}{4\pi}}.$$

Then we solve this algebraic equation

$$\hat{f}(\omega) = \frac{1}{1+i\omega-4\omega^2} e^{-\frac{\omega^2}{4\pi}}.$$

Finally, we take the inverse Fourier transform

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{1+i\omega-4\omega^2} e^{-\frac{\omega^2}{4\pi}} e^{i\omega x} d\omega.$$

1.MC4 [3 Points] The complex Fourier series of the function $\cosh(ax)$ on the interval $[-\pi, \pi]$ is given by

$$\sum_{n=-\infty}^{+\infty} \frac{(-1)^n a \sinh(a\pi)}{\pi(n^2 + a^2)} e^{inx}.$$

Find the value of the numerical series

$$\sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2}$$

(A) $\sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2} = \frac{\pi}{2a \sinh(a\pi)}.$

(B) $\sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2} = \frac{\pi}{2a \sinh(a\pi)} - \frac{1}{2a^2}.$

(C) $\sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2} = \frac{2\pi}{a \sinh(a\pi)} - \frac{2}{a^2}.$

(D) $\sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2} = \frac{2\pi}{a \sinh(a\pi)}.$

Solution:

The solution is (B) $\sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2} = \frac{\pi}{2a \sinh(a\pi)} - \frac{1}{2a^2}.$

The complex Fourier series will be equal to $\cosh(ax)$ for each $x \in [-\pi, \pi]$, or equivalently:

$$\frac{\pi \cosh(ax)}{a \sinh(a\pi)} = \sum_{n=-\infty}^{+\infty} \frac{(-1)^n}{n^2 + a^2} e^{inx}, \quad \forall x \in [-\pi, \pi]. \quad (1)$$

In particular for $x = 0$ we obtain something very similar to what we need. Observe that the right hand side becomes

$$\sum_{n=-\infty}^{+\infty} \frac{(-1)^n}{n^2 + a^2} \underbrace{e^{in0}}_{=1} = \sum_{n=-\infty}^{+\infty} \frac{(-1)^n}{n^2 + a^2} = \underbrace{\frac{1}{a^2}}_{\substack{n=0 \\ \text{term}}} + 2 \sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2} =: \frac{1}{a^2} + 2S,$$

where S is the sum we wanted to compute. Therefore equation (1) in $x = 0$ becomes:

$$\frac{\pi}{a \sinh(a\pi)} = \frac{1}{a^2} + 2S,$$

from which

$$S = \sum_{n=1}^{+\infty} \frac{(-1)^n}{n^2 + a^2} = \frac{\pi}{2a \sinh(a\pi)} - \frac{1}{2a^2}.$$

1.MC5 [3 Points] Determine if the following function is even, odd, or neither and if it is periodic or not. If the function is periodic, determine the fundamental period.

$$15 \cos(3x) + 3 \cos(4x).$$

- (A) The function is even and periodic of fundamental period 2π .
(B) The function is odd and periodic of fundamental period 2π .
(C) The function is even and periodic of fundamental period π .
(D) The function is odd and periodic of fundamental period π .

Solution:

(A) The function is even and periodic of fundamental period 2π .

First,

$$15 \cos(3(-x)) + 3 \cos(4(-x)) = 15 \cos(-3x) + 3 \cos(-4x) = 15 \cos(3x) + 3 \cos(4x).$$

Therefore, the function is even.

If $f(x)$ is periodic of period P_1 and $g(x)$ is periodic of period P_2 , then their sum $f(x) + g(x)$ is periodic of period the least common multiple $P = \text{LCM}(P_1, P_2)$ of the two periods.

In this case, $15 \cos(3x)$ is periodic of fundamental period $\frac{2\pi}{3}$ while $3 \cos(4x)$ is periodic of fundamental period $\frac{2\pi}{4}$, therefore their sum is periodic of period

$$P = \text{LCM}\left(\frac{2\pi}{3}, \frac{\pi}{2}\right) = \text{LCM}(4, 3) \cdot \frac{\pi}{6} = \frac{12}{6}\pi = 2\pi$$

It is easy to see that no smaller number is a period.

1.MC6 [3 Points] Consider the following PDE (partial differential equation) for the function $u = u(x, y)$:

$$4u_{xx} + xu_x + 6u_{xy} - yu_y = -7u_{yy} + u_x.$$

Is the PDE hyperbolic, parabolic, elliptic or of mixed type ?

- (A) hyperbolic.
(B) elliptic.
(C) parabolic.
(D) mixed type.

Solution:

(B) The PDE is elliptic because, $A = 4$, $B = 3$, and $C = 7$ therefore,

$$AC - B^2 = 4 \cdot 7 - 9 = 19 > 0.$$

The sign of the coefficient is positive, therefore the PDE is elliptic.

1.MC7 [3 Points] Wave equation with D'Alembert solution.

Let $u(x, t)$ be the solution of the following problem

$$\begin{cases} u_{tt} = u_{xx}, & x \in \mathbb{R}, t > 0, \\ u(x, 0) = f(x) = \begin{cases} 2, & |x| \leq 14 \\ 0, & |x| > 14 \end{cases} & x \in \mathbb{R}, \\ u_t(x, 0) = g(x) = \begin{cases} 1, & 3 \leq x \leq 4 \\ 0, & x \notin [3, 4] \end{cases} & x \in \mathbb{R}. \end{cases}$$

Find the values of u at the point $(x, t) = (10, 7)$, i.e. find $u(10, 7)$

- (A) $u(10, 7) = 8.$
- (B) $u(10, 7) = 10.$
- (C) $u(10, 7) = \frac{3}{2}.$
- (D) $u(10, 7) = \frac{5}{2}.$

Solution:

The solution is (C) $u(10, 7) = \frac{3}{2}.$

D'Alembert's formula for the solution of the wave equation is:

$$u(x, t) = \frac{1}{2} (f(x + ct) + f(x - ct)) + \frac{1}{2c} \int_{x-ct}^{x+ct} g(s) ds.$$

With our given initial conditions, $c = 1$, $x = 10$, and $t = 7$, we get

$$\begin{aligned} u(10, 7) &= \frac{1}{2} (f(17) + f(3)) + \frac{1}{2} \int_3^{17} g(s) ds \\ &= \frac{1}{2} (0 + 2) + \frac{1}{2} \int_3^4 1 ds \\ &= 1 + \frac{1}{2} \\ &= \frac{3}{2}. \end{aligned}$$

1.MC8 [3 Points] Let $u = u(x, y)$ be a harmonic function in D_2 the disk of radius 2 centred at 0.

The maximum value of u is at $(x, y) = (\sqrt{2}, -\sqrt{2})$, i.e. $\max_{D_2} u(x, y) = u(\sqrt{2}, -\sqrt{2})$.

Which of the following statements is true?

- (A) u is not constant in D_2 .
- (B) u is constant in D_2 .
- (C) There exists another point (x', y') in D_2 such that $u(x, y) = u(x', y')$.
- (D) We cannot conclude that (A), (B) and (C) are true for every u .

Solution:

The solution is (D): We cannot conclude that (A), (B) and (C) are true.

The point $(x, y) = (\sqrt{2}, -\sqrt{2})$ is on the boundary of D_2 .

Therefore, the maximum of u is taken on the boundary. Hence, we cannot conclude anything about u .

1.MC9 [3 Points] Consider the Dirichlet problem for the Laplace equation,

$$R = \{(x, y) \in \mathbb{R}^2 \mid 0 \leq x \leq 1, 0 \leq y \leq 2\},$$

$$\begin{cases} \Delta u = 0, & (x, y) \in R, \\ u(0, y) = u(1, y) = 0, & 0 \leq y \leq 2, \\ u(x, 0) = 0, & 0 \leq x \leq 1, \\ u(x, 2) = f(x) & 0 \leq x \leq 1, \end{cases}$$

where f is a continuous function.

- (A) There are infinitely many solutions.
- (B) We cannot say anything.
- (C) There is a unique solution.
- (D) There is no solution.

Solution:

(C) There is a unique solution.

By the lecture notes we know that there exists a solution. We can show that there is a unique solution in the following way:

Let assume that u_1 and u_2 are two solutions of the Laplace equation. And consider $v = u_1 - u_2$. Then by linearity of the Laplace equation we have,

$$\Delta v = \Delta u_1 - \Delta u_2 = 0.$$

And for the boundary conditions we have

$$\begin{aligned} v(0, y) &= u_1(0, y) - u_2(0, y) = 0, & 0 \leq y \leq 2, \\ v(1, y) &= u_1(1, y) - u_2(1, y) = 0, & 0 \leq y \leq 2, \\ v(x, 0) &= u_1(x, 0) - u_2(x, 0) = 0, & 0 \leq x \leq 1, \\ v(x, 2) &= u_1(x, 2) - u_2(x, 2) = f(x) - f(x) = 0 & 0 \leq x \leq 1. \end{aligned}$$

Therefore, we get for v ,

$$\begin{cases} \Delta v = 0, & (x, y) \in R, \\ v(0, y) = v(1, y) = 0, & 0 \leq y \leq 2, \\ v(x, 0) = 0, & 0 \leq x \leq 1, \\ v(x, 2) = 0 & 0 \leq x \leq 1. \end{cases}$$

Hence, $v = 0$ and $u_1 = u_2$.

Question 2

2.Q1 [15 Points] Heat Equation with inhomogeneous boundary conditions

Find the general solution of the Heat equation (**with inhomogeneous boundary conditions**) for the following problem:

$$(1) \quad \begin{cases} u_t = c^2 u_{xx}, & 0 \leq x \leq \pi, t \geq 0, \\ u(0, t) = 5, & t \geq 0, \\ u(\pi, t) = 8, & t \geq 0, \\ u(x, 0) = f(x) + w(x), & 0 \leq x \leq \pi, \end{cases}$$

where w is the function that you have to find in point a) and f is given by

$$f(x) = \begin{cases} x^2 & \text{if } x \in [0, \pi), \\ 0 & \text{if } x = \pi. \end{cases}$$

You must proceed as follows.

- Find the unique function $w = w(x)$ with $w'' = 0$, $w(0) = 5$, and $w(\pi) = 8$.
- Define $v(x, t) := u(x, t) - w(x)$. Formulate the corresponding problem for v , equivalent to (1).
- Find, using the formula from the script, the solution $v(x, t)$ of the problem you have just formulated.
- Write down explicitly the solution $u(x, t)$ of the original problem (1).

Solution:

- a) The only functions with second derivative zero are the linear functions

$$w(x) = \alpha x + \beta, \quad \alpha, \beta \in \mathbb{R}.$$

Imposing the boundary conditions we find the right coefficients

$$\begin{cases} 5 = w(0) = \alpha \cdot 0 + \beta \\ 8 = w(\pi) = \alpha \cdot \pi + \beta \end{cases} \Leftrightarrow \begin{cases} \alpha = \frac{8-5}{\pi} = \frac{3}{\pi} \\ \beta = 5 \end{cases} \Leftrightarrow w(x) = \frac{3}{\pi}x + 5.$$

- b) The PDE doesn't change because w is independent of time and has second derivative in x zero. The boundary conditions become homogeneous (that's why we chose this w)

$$v(0, t) = u(0, t) - w(0) = 5 - 5 = 0 \quad \text{and} \quad v(\pi, t) = u(\pi, t) - w(\pi) = 8 - 8 = 0.$$

The initial condition becomes

$$v(x, 0) = u(x, 0) - w(x) = f(x) + w(x) - w(x) = f(x).$$

Finally, the boundary value problem for v with homogeneous boundary conditions reads

as

$$\begin{cases} v_t = c^2 v_{xx}, & 0 \leq x \leq \pi, t \geq 0, \\ v(0, t) = 0, & t \geq 0, \\ v(\pi, t) = 0, & t \geq 0, \\ v(x, 0) = f(x) & 0 \leq x \leq \pi. \end{cases}$$

c) By the lecture notes the general solution of the heat equation in an interval is given by

$$v(x, t) = \sum_{n=1}^{\infty} A_n e^{-c^2 n^2 t} \sin(nx).$$

By imposing the initial condition $v(x, 0) = f(x)$, we have

$$v(x, 0) = \sum_{n=1}^{\infty} A_n \sin(nx) = f(x).$$

Hence the A_n are the Fourier coefficients of the extension of f to be an odd periodic function of period 2π , that is

$$\begin{aligned} A_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \sin(nx) dx = \frac{2}{\pi} \int_0^{\pi} x^2 \sin(nx) dx \\ &= \frac{2}{\pi} \frac{(2 - n^2 x^2) \cos(nx) + 2nx \sin(nx)}{n^3} \Big|_0^{\pi} \\ &= \frac{2}{\pi} \frac{(2 - n^2 \pi^2) \cos(n\pi)}{n^3} - \frac{2}{\pi} \frac{2}{n^3} \\ &= \frac{2}{\pi} \frac{(2 - n^2 \pi^2) (-1)^n}{n^3} - \frac{2}{\pi} \frac{2}{n^3} \\ &= \frac{4((-1)^n - 1)}{\pi n^3} - \frac{2\pi(-1)^n}{n}. \end{aligned}$$

Hence,

$$v(x, t) = \sum_{n=1}^{\infty} \left(\frac{4((-1)^n - 1)}{\pi n^3} - \frac{2\pi(-1)^n}{n} \right) e^{-c^2 n^2 t} \sin(nx).$$

d) The solution $u(x, t)$ is given by

$$u(x, t) = v(x, t) + w(x) = \sum_{n=1}^{\infty} \left(\frac{4((-1)^n - 1)}{\pi n^3} - \frac{2\pi(-1)^n}{n} \right) e^{-c^2 n^2 t} \sin(nx) + \frac{3}{\pi} x + 5.$$

Question 3

3.Q1 [10 Points] Dirichlet problem on a region with symmetries

Find the solution $u(r, \theta)$ of the following Dirichlet problem on the disk of radius R in polar coordinates:

$$\begin{cases} \Delta u = 0, & 0 \leq r \leq R, 0 \leq \theta \leq 2\pi, \\ u(R, \theta) = \sin^2(\theta) + 8 \cos^3(\theta), & 0 \leq \theta \leq 2\pi. \end{cases}$$

You should give the answer without unsolved integral and you can use the formulas developed in the lecture.

[Hint: Don't try to find the solution in the Poisson integral form.]

[Hint: Remember the trigonometric formulas

$$\sin^2(\theta) = \frac{1}{2} - \frac{1}{2} \cos(2\theta) \quad \text{and} \quad \cos^3(\theta) = \frac{3}{4} \cos(\theta) + \frac{1}{4} \cos(3\theta).]$$

Solution:

The solution will be

$$u(r, \theta) = \sum_{n=0}^{+\infty} r^n (A_n \cos(n\theta) + B_n \sin(n\theta)),$$

with coefficients found imposing

$$u(R, \theta) = \sum_{n=0}^{+\infty} R^n (A_n \cos(n\theta) + B_n \sin(n\theta)) = \sin^2(\theta) + 8 \cos^3(\theta).$$

Using the trigonometric formulas

$$\sin^2(\theta) = \frac{1}{2} - \frac{1}{2} \cos(2\theta) \quad \text{and} \quad 8 \cos^3(\theta) = 6 \cos(\theta) + 2 \cos(3\theta),$$

we have,

$$u(R, \theta) = \sum_{n=0}^{+\infty} R^n (A_n \cos(n\theta) + B_n \sin(n\theta)) = \frac{1}{2} + 6 \cos(\theta) - \frac{1}{2} \cos(2\theta) + 2 \cos(3\theta).$$

We obtain coefficients

$$\begin{cases} B_n = 0, & \forall n \geq 0, \\ A_n = 0, & \forall n \geq 0, n \neq 0, 1, 2, \text{ and } 3, \\ A_0 = \frac{1}{2}, \\ A_1 = \frac{6}{R}, \\ A_2 = -\frac{1}{2R^2}, \\ A_3 = \frac{2}{R^3}. \end{cases}$$

Finally

$$u(r, \theta) = \frac{1}{2} + \frac{6r}{R} \cos(\theta) - \frac{r^2}{2R^2} \cos(2\theta) + \frac{2r^3}{R^3} \cos(3\theta).$$