

Analysis Übungsstunde 5

qtuerler@student.ethz.ch

17.10.24



Komplexe Fourier Reihe

normale Fourier-Reihe: $a_0 + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi}{L}x\right) + b_n \sin\left(\frac{n\pi}{L}x\right)$

ersetze durch cplx Schreibweise $\sin(x) =$

$\cos(x) =$

$$f(x) = a_0 + \sum_{n=1}^{\infty} a_n \frac{1}{2}(e^{i\frac{n\pi}{L}x} + e^{-i\frac{n\pi}{L}x}) + b_n \frac{1}{2i}(e^{i\frac{n\pi}{L}x} - e^{-i\frac{n\pi}{L}x})$$

$\left[\frac{1}{i} = -i \right]$
 $\left(\frac{1}{i} \cdot \frac{1}{i} = \frac{1}{i^2} = -1 \right)$

$$= a_0 + \sum_{n=1}^{\infty} \frac{a_n - ib_n}{2} e^{i\frac{n\pi}{L}x} + \frac{a_n + ib_n}{2} e^{-i\frac{n\pi}{L}x}$$

$$= a_0 + \sum_{n=1}^{\infty} e^{i\frac{n\pi}{L}x} + e^{-i\frac{n\pi}{L}x}$$

$$= c_0 + \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} c_n \cdot e^{i\frac{n\pi}{L}x} \quad \text{oder} \quad \sum_{n=-\infty}^{\infty} c_n \cdot e^{i\frac{n\pi}{L}x}$$

(I) $\longleftrightarrow$ (II)

↙ passe einfach auf
dass $c_n \forall n \in \mathbb{Z}$ geht
bsp: $\frac{1}{n} \rightarrow c_0$ separat

$$f(x) = c_0 + \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} c_n \cdot e^{i\frac{\pi n}{L}x}$$

! Wie immer, $f(x)$ muss
2L-Periodisch sein!

$$c_n = \frac{1}{2L} \int_{-L}^L f(x) \cdot e^{-i\frac{\pi n}{L}x} dx ; \quad c_0 = \frac{1}{2L} \int_{-L}^L f(x) dx$$

$$a_0 = c_0 ; \quad a_n = c_n + c_{-n} ; \quad b_n = i(c_n - c_{-n})$$

$$e^{ix} + e^{-ix} = 2\cos(x); \quad e^{ix} - e^{-ix} = 2i\sin(x)$$

$$c_n = \frac{1}{2}(a_n - ib_n)$$

$$c_{-n} = \frac{1}{2}(a_n + ib_n)$$

Remembers:

$$\bullet e^{\pm ix} =$$

$$\bullet e^{i\pi} = \longrightarrow \bullet e^{\pm i n \pi} =$$


Bsp: Kplx Fourier-Reihe

Berechne Fourier Reihe von $f(x) = e^x$, $x \in [-\pi, \pi]$
mit Periode $f(x) = f(x + 2\pi)$

$$f(x) = c_0 + \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} c_n \cdot e^{\frac{i n \pi}{L} x}$$

$$\Rightarrow c_0 =$$

$$\Rightarrow c_n =$$



Bsp. Prüfung Sommer 2017 A2


2. Fourier Series (6+1=7 Points)

- a) Given the odd, 2π -periodic function $f: \mathbb{R} \rightarrow \mathbb{R}$ with

$$f(x) = \begin{cases} x & 0 \leq x \leq \frac{\pi}{2} \\ -(x - \pi) & \frac{\pi}{2} \leq x \leq \pi, \end{cases}$$

determine the real Fourier series of f , i.e. state it's Fourier series and determine the coefficients.

- b) Does the Fourier series converge pointwise to the function f ? Please justify your answer.



$\Rightarrow$

$\Rightarrow$

$b_0 =$

$$\begin{aligned} \int x \sin(nx) &= -\frac{x}{n} \cos(nx) + \frac{1}{n} \int \cos(nx) dx \\ &= -\frac{x}{n} \cos(nx) + \frac{1}{n^2} \sin(nx) = \left[\frac{\sin(nx) - nx \cos(nx)}{n^2} \right] \end{aligned}$$

$\leftarrow$ Grenzen einsetzen

$$\left[\begin{array}{l} \text{FALLS:} \\ \text{(hypothetisch)} \end{array} \right. = \left\{ \begin{array}{ll} \frac{1}{k} & n=2k \\ (-1)^k & n=2k+1 \end{array} \right. \quad L=\pi$$

$$f(x) =$$

Alle geraden

ungeraden Terme

Für Serie: Brauche auch Skript letzte Woche!